

AUTHOR INDEX

Bold type indicates authors of commentaries and editorials; (No.) indicates issue number; (L) indicates letter to the editor.

- A**
 Abrahamowicz M, 12
 AIDS Clinical Trials Group 320 Study Team, 43
 American Society of Addiction Medicine Working Group on Pharmacological Management of Alcohol Withdrawal, 8
 Andersen HR, 62
 Angioplasty Substudy Investigators, 6
 Antiarrhythmics Versus Implantable Defibrillators (AVID) Investigators, 60
Arnold D, 72
Atallah AN, 63
Atkins D, A-12-14(No. 2)
 Avins AL, 10
- B**
 Badgett RG, 11
 Bakke F, 29
 Balas EA, 22
 Bartalena L, 71
 Battista RN, 12
Bauer DC, 47
 Bayerdörffer E, 19
 Beck E, 35
 Beech R, 58
Beghe C, 81
 Berkowitz SD, 17
 Berman N, 70
 Beshansky JR, 25
 Betriu A, 59
 Bigger JT Jr, 61
 Blauw GJ, 76
 Blum AL, 19
 Bogazzi F, 71
Boissel JP, 35
 Boivin JF, 15
Bostom AG, 21
 Bouges S, 5
 Bouter LM, 65
 Boyle DI, 50
 Bozovich G, 35
Bradley K, 51
 Britton M, 81
Bronson DL, 64
 Browner WS, 10
Brun-Boisson C, 40-41
 Brunner E, 37
Buntinx F, 64, 78
 Burgess E, 63
 Byington RP, 1
 Byttemier G, 64
- C**
Cairns JA, 60-61
Calam J, 19
 Canadian Coordinating Office for Health Technology Assessment (CCOHTA), 46
Carney T, 57
 Caro J, 80
 Cerhan JR, 74
- Chan FK, 44
 Chaulk P, 24
Chick J, 8
 Chung SC, 44
 Clapperton M, 32
Clarke R, 38, 76
Clutter WE, 71
 Cohen M, 34
 Colditz GA, 14
 Collaborative Group on Hormonal Factors in Breast Cancer, 73
 Cook DJ, 39
 Coronary Artery Bypass Graft (CABG) Patch Trial Investigators, 61
 Côté R, 12
 Crouse JR 3d, 1
 Cumming R, 20, 69
Cummings SR, 49
- D**
 Dallal GE, 47
 Daly LE, 21
 Daroca A, 35
 Darouiche R, 41
 DARTS/MEMO Collaboration, 50
 Davey Smith G, 31
 Dawson-Hughes B, 47
 Deeks J, 45
 Demers C, 34
 Demicheli V, 45
Devins GM, 10
 Diem P, 31
 Dinneen SF, 16
 Dupuis J, 41
- E**
Easterbrook P, 13
 Efficacy and Safety of Subcutaneous Enoxaparin in Non-Q-Wave Coronary Events Study Group, 34
 Egger M, 31
Ehrlich GE, 65
Eisenberg JM, A-12-14(No. 2)
Eisenberg PR, 4, 17
 Ernst E, 38
Eslick GD, 44
 Evans DJ, 66
- F**
Faris I, 77
Feagan B, 33
Feinmann C, 3
 Fick G, 33
 Fine MJ, 72
Fletcher SW, A-14-16(No. 3)
 Formoterol and Corticosteroids Establishing Therapy (FACET) International Study Group, 67
 Freaake D, 24
 Funnell MM, 30
 Furberg CD, 1
- G**
Geerts WH, 5
 Gerstein HC, 16
 Gillespie LD, 69
 Gillespie WJ, 69
 Giorgi JV, 13
 Glennie J, 46
 Global Use of Strategies to Open Occluded Coronary Arteries in Acute Coronary Syndromes (GUSTO IIb), 6
 Global Utilization of Streptokinase and Tissue Plasminogen Activator for Occluded Coronary Arteries (GUSTO) I Investigators, 17
 Goldberg DE, 51
 Gotzsche PC, 9
 Gough A, 32
 Graham IM, 21
 Grande P, 4
 Granger CB, 17
Gray A, 80
 Griffin S, 57
 Griffith JL, 25
 Grodstein F, 14
 Gulick RM, 42
 Gurfinkel E, 34, 35
- H**
 Hackshaw AK, 49
 Häheim LL, 29
 Hammer SM, 43
 Harris SS, 47
Harvey RL, 29, 58
 Havlir D, 42
 Haynes RB, 23
 Hemmelgarn B, 15
 Hemminki E, 48
Hensley MJ, 20, 72
 Herman WH, 30
Hirsh CH, 46
Hirsh J, 52
 Hoen HM, 1
Holbrook A, 15
Holden JD, 15
Holloway R, 12
 Hopper AH, 81
 Hu FB, 75
 Huang A, 15
Hughes I, 15
 Hughes MD, 43
- I**
 Indredavik B, 29
- J**
 Jaffrey F, 22
Jamal SA, 49
 Jefferson T, 45
Johnson BE, 74
Jones TV, 69
 Jung AC, 3

- K**
Kamerow D, A-12-14(No. 2)
Kaplan GA, 51
Kapoor WN, 72
Katz MM, 70
Kauhanen J, 51
Keenan SP, 39
Kerlikowske K, 79
Kernerman PD, 39
Kinmonth AL, 57
Klittich W, 80
Koch A, 5
Koes BW, 65
Krahn M, 45
Krall EA, 47
Kuperman GJ, 22
- L**
Labenz J, 19
Lagaay AM, 76
Lamb SE, 69
Lange IR, 63
Law MR, 49
Le Bars PL, 70
Leduc L, 63
Leeder SR, 20
Leiter LA, 30
LeLorier J, 63
Lewis RF, 12
Lip GYH, 62
Löfdahl CG, 67
Lonn EM, 7
Lucy CR, 11
Luckmann R, 70
- M**
Madsen JK, 4
Maki DG, 40
Malenka DJ, 36
Manson JE, 75
Mancocci C, 71
Martin FC, 81
Massel D, 59
Mautner B, 35
May GR, 33
Mayo-Smith MF, 8
McDermott MM, 53
McDonald CJ, 22
McGinn T, A-14-15(No.1), 25
McGuire A, 80
McKibbin KA, A-15(No. 2)
McMahon AD, 50
McNagny SE, 14
McPherson K, 48
Mellors JW, 13, 42
Mensah EK, 78
Mermal LA, 40
Metlay JP, 72
Middleman EL, 23
Mink PJ, 74
Miranda J, 10
- Mitchell P, 20
Morris AD, 50
Morris AH, 39
Mullen PD, 68
Mulrow CD, 11
Muñoz A, 13
Mushlin A, 79
- N**
Ness RB, 48
Niedzwiecki D, 2
Nielsen JC, 62
North American EGb Study Group, 70
- O**
Ohman EM, 34
Oliver D, 81
Oxman AD, 23
- P**
Pauwels RA, 67
Pearson DC, 33
Peterson WL, 9
Phillips K, 79
Pieper KS, 17
Pinard G, 15
Postma DS, 67
Price DT, 52
Pugh JA, 50
- R**
Raad I, 41
Ramírez G, 68
Randolph A, A-14-15(No. 1)
Rapola JM, 7
Redfern C, 1
Refsum HM, 21
Rembold C, 2
Rey E, 63
Richardson S, A-14-15(No. 1)
Ridker PM, 52
Ripatti S, 7
RITA-2 Trial Participants, 36
Roberts RR, 78
Rokseth R, 29
Rolando N, 32
Rothwell PM, 77
Rowe BH, 69
ROXIS Study Group, 35
Rudd AG, 58
Rudd P, 53
- S**
Sackett D, A-14-15(No. 1)
Salonen JT, 51
Salzmann P, 79
Saunamäki K, 4
Schmid CH, 2
Schmitt B, 6, 53
Seed P, 81
Selker HP, 25
- Sellers TA, 74
Sherman SE, 68
Silagy C, 18
Simoes RJ, 59
Simon JA, 37, 75
Simons-Morton DG, 68
Slattery J, 77
Slørdahl SA, 29
Smith BJ, 66-67
Smith DA, 16
Squires KE, 43
Staessen JA, 64
Staiger T, 3
Stampfer MJ, 14, 75
Standiford CJ, 30
Stason WB, 2
Stettler C, 31
Stevens DL, 30
Stolz SM, 40
Suissa S, 15
Sullivan M, 3
Sung JJ, 44
Sutherland LR, 33
Szerlip HM, 11
- T**
Talley NJ, 44
Taylor DA, 66
Thomsen PE, 62
Thomson MA, 23
Thorogood M, 37
Tilling K, 58
- V**
van Tulder MW, 65
Vijan S, 30
Virtamo J, 7
- W**
Wald NJ, 18
Waldman MH, 32
Wallner E, 53
Warlow CP, 77
Watt HC, 18
Weaver WD, 59
Weverling-Rijnsburger AW, 76
Wheeler S, 40
White I, 37
Whooley MA, 10
Williams CJ, 73
Wilson MC, 24
Wolfe CD, 58
Worrall G, 24
- Z**
Zalenski RJ, 78
Zar FA, 42-43
Zetterstrom O, 66
Ziegler S, 5
Zinman B, 31

SUBJECT INDEX

A

- Accidents, traffic
 - Use of long-half-life benzodiazepines in older persons was associated with increased risk for injurious motor vehicle accidents, 15
- Adrenal cortex hormones
 - Use of inhaled corticosteroids was associated with the development of cataracts, 20
- Aged
 - Calcium and vitamin D reduced nonvertebral fractures and bone loss, 47
 - High total cholesterol levels reduced the risk for all-cause mortality in adults older than 85 years of age, 76
 - Review: Multiple risk factor modification reduces falls in elderly persons, 69
 - A 6-point risk score predicted which elderly patients would fall in hospital, 81
 - Use of long-half-life benzodiazepines in older persons was associated with increased risk for injurious motor vehicle accidents, 15
- AIDS (*see also* HIV)
 - HIV-1 RNA levels and CD4⁺ cell counts predicted disease progression in HIV infection, 13
- Albuminuria
 - Review: Microalbuminuria predicts overall mortality and cardiovascular mortality and morbidity in non-insulin-dependent diabetes mellitus, 16
- Alcohol drinking
 - Binge beer drinking was associated with mortality in middle-aged men, 51
 - Review: Benzodiazepines are more effective than neuroleptics in reducing delirium and seizures in alcohol withdrawal, 8
- Alcohol withdrawal delirium
 - Review: Benzodiazepines are more effective than neuroleptics in reducing delirium and seizures in alcohol withdrawal, 8
- Alteplase
 - Older age, lighter weight, female sex, African ancestry, and invasive procedures were associated with bleeding after thrombolytic therapy, 17
 - Primary angioplasty was marginally better than t-PA after MI, 6
- Alzheimer disease
 - Ginkgo biloba safely and modestly improved dementia, 70
 - Review: Tacrine has limited efficacy for Alzheimer disease, 46
- Angina pectoris
 - Enoxaparin reduced combined coronary events in unstable angina and non-Q-wave MI at 14 and 30 days, 34
 - In a pilot study, roxithromycin reduced 1 of 10 end points of ischemic cardiac disease, 35
- Angioplasty, transluminal, percutaneous coronary
 - Coronary angioplasty led to more nonfatal MIs than medical therapy in patients with coronary artery disease, 36
 - Invasive treatment reduced reinfarctions and admissions for unstable angina in patients with ischemia after MI, 4

- Primary angioplasty was marginally better than t-PA after MI, 6
- Review: Early angioplasty improves outcomes for patients with myocardial infarction, 59
- Angiotensin-converting enzyme inhibitors
 - ACE inhibitor use was associated with severe hypoglycemia in patients with diabetes mellitus, 50
- Antiarrhythmics Versus Implantable Defibrillators (AVID) Trial
 - Implantable cardioverter-defibrillators reduced mortality in patients resuscitated from ventricular arrhythmias, 60
- Antibiotics, macrolide
 - In a pilot study, roxithromycin reduced 1 of 10 end points of ischemic cardiac disease, 35
- Antihypertensive agents
 - Ambulatory blood pressure monitoring allowed more patients to stop antihypertensive drug therapy, 64
 - Review: Pharmacologic treatment of hypertension during pregnancy reduces perinatal deaths and severe maternal hypertension, 63
- Arterial occlusive diseases
 - Review: Chelation therapy is ineffective for peripheral arterial occlusive disease, 38
- Asthma
 - Formoterol and budesonide reduced exacerbations and improved lung function in patients with stable asthma, 67
 - Low-dose oral theophylline plus budesonide was as effective as high-dose budesonide for treatment of moderate asthma, 66
- Atrial fibrillation
 - Atrial pacing was better than ventricular pacing for the sick-sinus syndrome, 62
 - Implantable cardioverter-defibrillators reduced mortality in patients resuscitated from ventricular arrhythmias, 60
- Azathioprine
 - Review: Azathioprine maintains remission in Crohn disease, but withdrawal rates are higher, 33

B

- β -carotene
 - Vitamin E and β -carotene were not effective in men at high risk for coronary events, 7
- Beclomethasone
 - Use of inhaled corticosteroids was associated with the development of cataracts, 20
- Beer
 - Binge beer drinking was associated with mortality in middle-aged men, 51
- Benzodiazepines
 - Review: Benzodiazepines are more effective than neuroleptics in reducing delirium and seizures in alcohol withdrawal, 8
 - Use of long-half-life benzodiazepines in older persons was associated with increased risk for injurious motor vehicle accidents, 15
- Blood pressure monitoring
 - Ambulatory blood pressure monitoring allowed more patients to stop antihypertensive drug therapy, 64

Bone density

Calcium and vitamin D reduced nonvertebral fractures and bone loss, 47

Review: Smoking is associated with increased hip fracture in postmenopausal women, 49

Breast neoplasms

Extending screening mammography to include younger women minimally increased life expectancy with large increases in cost, 79

HRT was not associated with increased breast cancer in women with a positive family history of breast cancer, 74

Review: Recent HRT increases the risk for breast cancer, 73

Bronchodilator agents

Formoterol and budesonide reduced exacerbations and improved lung function in patients with stable asthma, 67

Low-dose oral theophylline plus budesonide was as effective as high-dose budesonide for treatment of moderate asthma, 66

Budesonide

Formoterol and budesonide reduced exacerbations and improved lung function in patients with stable asthma, 67

Low-dose oral theophylline plus budesonide was as effective as high-dose budesonide for treatment of moderate asthma, 66

C

Calcium

Calcium and vitamin D reduced nonvertebral fractures and bone loss, 47

Cardiac pacing

Atrial pacing was better than ventricular pacing for the sick-sinus syndrome, 62

Cardiovascular diseases

High total cholesterol levels reduced the risk for all-cause mortality in adults older than 85 years of age, 76

Pravastatin cost was £20 375 (U.S. \$34 360) per life-year gained for primary prevention among men with increased risk for cardiovascular disease, 80

Review: Dietary advice modestly reduces blood pressure, serum cholesterol, and urinary sodium levels, 37

Review: HRT is not associated with cardiovascular events or cancer in postmenopausal women in recent clinical trials, 48

Review: Microalbuminuria predicts overall mortality and cardiovascular mortality and morbidity in non-insulin-dependent diabetes mellitus, 16

Carotid stenosis

Serial ultrasonography did not predict stroke in asymptomatic carotid artery disease, 12

Cataract

Use of inhaled corticosteroids was associated with the development of cataracts, 20

Cellulitis

Granulocyte-colony stimulating factor hastened healing of severe foot infections in patients with diabetes mellitus, 32

Central venous catheterization

Antiseptic catheters reduced catheter-related infections and were well tolerated, 40

A catheter coated with minocycline and rifampin reduced bloodstream infections, 41

Cerebrovascular disorders

Early discharge to community care was as effective as conventional care for stroke, 58

Review: Statins prevent stroke, especially in patients with coronary heart disease, 1

Review: Stroke and death from carotid endarterectomy are predicted by 5 clinical and 3 angiographic factors, 77

Serial ultrasonography did not predict stroke in asymptomatic carotid artery disease, 12

Stroke unit care improved survival and function for 5 years after an acute stroke, 29

Chelation therapy

Review: Chelation therapy is ineffective for peripheral arterial occlusive disease, 38

Chest pain

An accelerated emergency department diagnostic protocol for chest pain reduced hospitalization and costs, 78

Chlorhexidine

Antiseptic catheters reduced catheter-related infections and were well tolerated, 40

Cholesterol

High total cholesterol levels reduced the risk for all-cause mortality in adults older than 85 years of age, 76

Citalopram

Review: Selective serotonin reuptake inhibitors are effective for mixed chronic pain, 3

Clinical competence

Review: Subtle clinical findings can detect left-sided heart failure in adults, 11

Clinical protocols

An accelerated emergency department diagnostic protocol for chest pain reduced hospitalization and costs, 78

Cognition

Ginkgo biloba safely and modestly improved dementia, 70

Review: Tacrine has limited efficacy for Alzheimer disease, 46

Communication

Review: Telephone and computer communication improves clinical health care process and patient outcome, 22

Community health services

Early discharge to community care was as effective as conventional care for stroke, 58

Community-acquired infections

Review: Clinical assessment is inaccurate for diagnosing community-acquired pneumonia, 72

Complications

Review: Several interventions reduce complications in type 2 diabetes mellitus, 30

Computers

Review: Telephone and computer communication improves clinical health care process and patient outcome, 22

Contraceptives, oral, adverse effects

- Review: Factor V Leiden mutation increases risk for thromboembolism, 52
- Coronary artery bypass graft surgery
 - Implantable cardioverter-defibrillators used at the time of bypass surgery did not improve survival among high-risk patients, 61
 - Invasive treatment reduced reinfarctions and admissions for unstable angina in patients with ischemia after MI, 4
- Coronary disease
 - Coronary angioplasty led to more nonfatal MIs than medical therapy in patients with coronary artery disease, 36
 - Dietary intake of monounsaturated and polyunsaturated fats reduced risk for coronary disease in women, 75
 - Review: Medication adherence may reduce mortality and morbidity, 53
 - Review: Statins prevent stroke, especially in patients with coronary heart disease, 1
- Counseling
 - Review: Patient education and counseling increase preventive behaviors, 68
- Crohn disease
 - Review: Azathioprine maintains remission in Crohn disease, but withdrawal rates are higher, 33

D

- Defibrillators, implantable
 - Implantable cardioverter-defibrillators reduced mortality in patients resuscitated from ventricular arrhythmias, 60
 - Implantable cardioverter-defibrillators used at the time of bypass surgery did not improve survival among high-risk patients, 61
- Dementia
 - Ginkgo biloba safely and modestly improved dementia, 70
- Depression
 - 2 questions were as good as more questions for detecting depression, 10
- Diabetes mellitus
 - ACE inhibitor use was associated with severe hypoglycemia in patients with diabetes mellitus, 50
 - Granulocyte-colony stimulating factor hastened healing of severe foot infections in patients with diabetes mellitus, 32
 - Review: Intensive insulin therapy for type 1 diabetes mellitus increases risk for severe hypoglycemia, and pump therapy increases ketoacidosis, 31
 - Review: Microalbuminuria predicts overall mortality and cardiovascular mortality and morbidity in non-insulin-dependent diabetes mellitus, 16
 - Review: Prompted follow-up by specialists and GPs produces similar outcomes for diabetes mellitus, 57
 - Review: Several interventions reduce complications in type 2 diabetes mellitus, 30
- Diabetic complications
 - Review: Intensive insulin therapy for type 1 diabetes mellitus increases risk for severe hypoglycemia, and pump therapy increases ketoacidosis, 31
 - Review: Several interventions reduce complications in type 2 diabetes mellitus, 30

Diet

- Dietary intake of monounsaturated and polyunsaturated fats reduced risk for coronary disease in women, 75
- Review: Dietary advice modestly reduces blood pressure, serum cholesterol, and urinary sodium levels, 37
- Drug effects
 - Review: Tacrine has limited efficacy for Alzheimer disease, 46
- Duodenal ulcer
 - Cure of *Helicobacter pylori* infection was associated with reflux esophagitis, 19

E

Editorial

- Clinical prediction guides, A-14-15(No.1)
- Evidence-based medicine at the Agency for Health Care Policy and Research, A-12-14(No. 2)
- Evidence-based screening: What kind of evidence is needed?, A-14(No. 3)
- Using *Best Evidence* in clinical practice, A-15(No. 2)
- Education, continuing
 - Review: Effectiveness of local opinion leaders varies, 23
- Education, patient
 - Review: Patient education and counseling increase preventive behaviors, 68
- Emergency services, hospital
 - An accelerated emergency department diagnostic protocol for chest pain reduced hospitalization and costs, 78
 - Benefit from thrombolysis in acute MI was predicted in the emergency department, 25
 - Review: Effectiveness of local opinion leaders varies, 23
- Endarterectomy, carotid
 - Review: Stroke and death from carotid endarterectomy are predicted by 5 clinical and 3 angiographic factors, 77
- Enoxaparin
 - Enoxaparin reduced combined coronary events in unstable angina and non-Q-wave MI at 14 and 30 days, 34
- Esophagitis, peptic
 - Cure of *Helicobacter pylori* infection was associated with reflux esophagitis, 19
- Estrogen replacement therapy
 - HRT was not associated with increased breast cancer in women with a positive family history of breast cancer, 74
 - Postmenopausal hormone use was associated with decreased total mortality, 14
 - Review: Factor V Leiden mutation increases risk for thromboembolism, 52
 - Review: HRT is not associated with cardiovascular events or cancer in postmenopausal women in recent clinical trials, 48
 - Review: Recent HRT increases the risk for breast cancer, 73

F

Factor V

- Review: Factor V Leiden mutation increases risk for thromboembolism, 52

Falls

Review: Multiple risk factor modification reduces falls in elderly persons, 69

A 6-point risk score predicted which elderly patients would fall in hospital, 81

Fluoxetine

Review: Selective serotonin reuptake inhibitors are effective for mixed chronic pain, 3

Fluvoxamine

Review: Selective serotonin reuptake inhibitors are effective for mixed chronic pain, 3

Forecasting

Benefit from thrombolysis in acute MI was predicted in the emergency department, 25

Review: Effectiveness of local opinion leaders varies, 23

Fractures

Calcium and vitamin D reduced nonvertebral fractures and bone loss, 47

Review: Smoking is associated with increased hip fracture in postmenopausal women, 49

G

Genetics

Review: Factor V Leiden mutation increases risk for thromboembolism, 52

Global Use of Strategies to Open Occluded Coronary Arteries in Acute Coronary Syndromes [GUSTO IIb]

Primary angioplasty was marginally better than t-PA after MI, 6

Glucocorticoids

Low-dose oral theophylline plus budesonide was as effective as high-dose budesonide for treatment of moderate asthma, 66

Formoterol and budesonide reduced exacerbations and improved lung function in patients with stable asthma, 67

Granulocyte-colony stimulating factor

Granulocyte-colony stimulating factor hastened healing of severe foot infections in patients with diabetes mellitus, 32

Graves disease

Radiiodine plus prednisone prevented development or progression of ophthalmopathy in patients with Graves hyperthyroidism, 71

H

Health personnel

Review: Plasma-derived hepatitis B vaccine is safe and effective for health care workers, 45

Heart failure, congestive

Review: Medication adherence may reduce mortality and morbidity, 53

Helicobacter pylori

Cure of *Helicobacter pylori* infection was associated with reflux esophagitis, 19

Eradication of *Helicobacter pylori* before NSAID treatment prevented ulcers, 44

Hematoma

Review: Low-molecular-weight heparin is at least as effective as unfractionated heparin for preventing thrombosis and embolism after surgery, 5

Hemorrhage

Older age, lighter weight, female sex, African ancestry, and invasive procedures were associated with bleeding after thrombolytic therapy, 17

Review: Somatostatin or octreotide reduces transfusion needs but not other outcomes in acute bleeding esophageal varices, 9

Heparin

Enoxaparin reduced combined coronary events in unstable angina and non-Q-wave MI at 14 and 30 days, 34

Older age, lighter weight, female sex, African ancestry, and invasive procedures were associated with bleeding after thrombolytic therapy, 17

Primary angioplasty was marginally better than t-PA after MI, 6

Review: Low-molecular-weight heparin is at least as effective as unfractionated heparin for preventing thrombosis and embolism after surgery, 5

Hepatitis B vaccines

Review: Plasma-derived hepatitis B vaccine is safe and effective for health care workers, 45

Hirudin

Primary angioplasty was marginally better than t-PA after MI, 6

HIV

Indinavir plus zidovudine and lamivudine reduced HIV RNA levels in most HIV-infected persons with previous antiretroviral treatment, 42

Indinavir plus zidovudine and lamivudine reduced the progression to AIDS in patients with previous zidovudine therapy, 43

Homocysteine

Elevated total homocysteine levels increased the risk for vascular disease, 21

Hospital costs

An accelerated emergency department diagnostic protocol for chest pain reduced hospitalization and costs, 78

Hospitalization

ACE inhibitor use was associated with severe hypoglycemia in patients with diabetes mellitus, 50

Review: Medication adherence may reduce mortality and morbidity, 53

A 6-point risk score predicted which elderly patients would fall in hospital, 81

Stroke unit care improved survival and function for 5 years after an acute stroke, 29

Hypercholesterolemia

Pravastatin cost was £20 375 (U.S. \$34 360) per life-year gained for primary prevention among men with increased risk for cardiovascular disease, 80

Hypertension

- Ambulatory blood pressure monitoring allowed more patients to stop antihypertensive drug therapy, 64
- Review: Nifedipine for hypertension may not increase the risk for adverse cardiovascular events, 2
- Review: Pharmacologic treatment of hypertension during pregnancy reduces perinatal deaths and severe maternal hypertension, 63

Hypoglycemia

- ACE inhibitor use was associated with severe hypoglycemia in patients with diabetes mellitus, 50
- Review: Intensive insulin therapy for type 1 diabetes mellitus increases risk for severe hypoglycemia, and pump therapy increases ketoacidosis, 31

I

Insulin

- Review: Intensive insulin therapy for type 1 diabetes mellitus increases risk for severe hypoglycemia, and pump therapy increases ketoacidosis, 31

Iowa Women's Health Study

- HRT was not associated with increased breast cancer in women with a positive family history of breast cancer, 74

L

Leadership

- Review: Effectiveness of local opinion leaders varies, 23

Length of stay

- Granulocyte-colony stimulating factor hastened healing of severe foot infections in patients with diabetes mellitus, 22

Lovastatin

- Review: Statins prevent stroke, especially in patients with coronary heart disease, 1

Low-back pain

- Review: NSAIDs and muscle relaxants reduce acute low-back pain; manipulation, back schools, and exercise reduce chronic low-back pain, 65

Lung diseases, obstructive

- Switching from cigarettes to pipes or cigars reduced smoking-related mortality, 18

Lung neoplasms

- Switching from cigarettes to pipes or cigars reduced smoking-related mortality, 18

M

Mammography

- Extending screening mammography to include younger women minimally increased life expectancy with large increases in cost, 79

Mass screening

- Extending screening mammography to include younger women minimally increased life expectancy with large increases in cost, 79

Medical history taking

- Review: Clinical assessment is inaccurate for diagnosing community-acquired pneumonia, 72

Men

- Binge beer drinking was associated with mortality in middle-aged men, 51
- HIV-1 RNA levels and CD4⁺ cell counts predicted disease progression in HIV infection, 13
- Pravastatin cost was £20 375 (U.S. \$34 360) per life-year gained for primary prevention among men with increased risk for cardiovascular disease, 80
- Vitamin E and β -carotene were not effective in men at high risk for coronary events, 7

Menopause

- Review: Smoking is associated with increased hip fracture in postmenopausal women, 49

Mercaptopurine

- Review: Azathioprine maintains remission in Crohn disease, but withdrawal rates are higher, 33

Methimazole

- Radiiodine plus prednisone prevented development or progression of ophthalmopathy in patients with Graves hyperthyroidism, 71

Minocycline

- A catheter coated with minocycline and rifampin reduced bloodstream infections, 41

Morbidity

- Review: Medication adherence may reduce mortality and morbidity, 53

Mortality

- Binge beer drinking was associated with mortality in middle-aged men, 51
- Coronary angioplasty led to more nonfatal MIs than medical therapy in patients with coronary artery disease, 36
- Enoxaparin reduced combined coronary events in unstable angina and non-Q-wave MI at 14 and 30 days, 34
- High total cholesterol levels reduced the risk for all-cause mortality in adults older than 85 years of age, 76
- HIV-1 RNA levels and CD4⁺ cell counts predicted disease progression in HIV infection, 13
- Implantable cardioverter-defibrillators reduced mortality in patients resuscitated from ventricular arrhythmias, 60
- Implantable cardioverter-defibrillators used at the time of bypass surgery did not improve survival among high-risk patients, 61
- Postmenopausal hormone use was associated with decreased total mortality, 14
- Review: Medication adherence may reduce mortality and morbidity, 53
- Review: Microalbuminuria predicts overall mortality and cardiovascular mortality and morbidity in non-insulin-dependent diabetes mellitus, 16
- Review: Noninvasive positive-pressure ventilation reduces mortality and need for intubation in the intensive care unit, 39
- Review: Somatostatin or octreotide reduces transfusion needs but not other outcomes in acute bleeding esophageal varices, 9
- Review: Stroke and death from carotid endarterectomy are predicted by 5 clinical and 3 angiographic factors, 77

Stroke unit care improved survival and function for 5 years after an acute stroke, 29

Vitamin E and β -carotene were not effective in men at high risk for coronary events, 7

Myocardial infarction

Benefit from thrombolysis in acute MI was predicted in the emergency department, 25

Coronary angioplasty led to more nonfatal MIs than medical therapy in patients with coronary artery disease, 36

Enoxaparin reduced combined coronary events in unstable angina and non-Q-wave MI at 14 and 30 days, 34

In a pilot study, roxithromycin reduced 1 of 10 end points of ischemic cardiac disease, 35

Older age, lighter weight, female sex, African ancestry, and invasive procedures were associated with bleeding after thrombolytic therapy, 17

Primary angioplasty was marginally better than t-PA after MI, 6

Review: Early angioplasty improves outcomes for patients with myocardial infarction, 59

Review: Effectiveness of local opinion leaders varies, 23

Vitamin E and β -carotene were not effective in men at high risk for coronary events, 7

Myocardial ischemia

Switching from cigarettes to pipes or cigars reduced smoking-related mortality, 18

N

Nebulizers and vaporizers

Use of inhaled corticosteroids was associated with the development of cataracts, 20

Neoplasms

Review: HRT is not associated with cardiovascular events or cancer in postmenopausal women in recent clinical trials, 48

Nifedipine

Review: Nifedipine for hypertension may not increase the risk for adverse cardiovascular events, 2

Nonsteroidal anti-inflammatory agents

Eradication of *Helicobacter pylori* before NSAID treatment prevented ulcers, 44

Review: NSAIDs and muscle relaxants reduce acute low-back pain; manipulation, back schools, and exercise reduce chronic low-back pain, 65

Nurses' Health Study

Postmenopausal hormone use was associated with decreased total mortality, 14

O

Occupational diseases, prevention and control

Review: Plasma-derived hepatitis B vaccine is safe and effective for health care workers, 45

Outcome and process assessment

An accelerated emergency department diagnostic protocol for chest pain reduced hospitalization and costs, 78

P

Pain

Review: Selective serotonin reuptake inhibitors are effective for mixed chronic pain, 3

Paroxetine

Review: Selective serotonin reuptake inhibitors are effective for mixed chronic pain, 3

Patient compliance

Review: Medication adherence may reduce mortality and morbidity, 53

Patient discharge

Early discharge to community care was as effective as conventional care for stroke, 58

Patient outcomes

Review: Clinical practice guidelines may not improve patient outcomes in primary care, 24

Peptic ulcer

Eradication of *Helicobacter pylori* before NSAID treatment prevented ulcers, 44

Physical examination

Review: Clinical assessment is inaccurate for diagnosing community-acquired pneumonia, 72

Review: Subtle clinical findings can detect left-sided heart failure in adults, 11

Physical therapy

Review: NSAIDs and muscle relaxants reduce acute low-back pain; manipulation, back schools, and exercise reduce chronic low-back pain, 65

Physicians' practice patterns

Review: Effectiveness of local opinion leaders varies, 23

Plant extracts

Ginkgo biloba safely and modestly improved dementia, 70

Pneumonia

Review: Clinical assessment is inaccurate for diagnosing community-acquired pneumonia, 72

Point mutation

Review: Factor V Leiden mutation increases risk for thromboembolism, 52

Practice guideline

Review: Benzodiazepines are more effective than neuroleptics in reducing delirium and seizures in alcohol withdrawal, 8

Review: Clinical practice guidelines may not improve patient outcomes in primary care, 24

Pravastatin

Pravastatin cost was \$20 375 (U.S. \$34 360) per life-year gained for primary prevention among men with increased risk for cardiovascular disease, 80

Review: Statins prevent stroke, especially in patients with coronary heart disease, 1

Prednisone

Radioiodine plus prednisone prevented development or progression of ophthalmopathy in patients with Graves hyperthyroidism, 71

Pregnancy

Review: Pharmacologic treatment of hypertension during pregnancy reduces perinatal deaths and severe maternal hypertension, 63

Prevention and control

Review: Dietary advice modestly reduces blood pressure, serum cholesterol, and urinary sodium levels, 37

Review: Several interventions reduce complications in type 2 diabetes mellitus, 30

Primary care

Review: Clinical practice guidelines may not improve patient outcomes in primary care, 24

Review: Prompted follow-up by specialists and GPs produces similar outcomes for diabetes mellitus, 57
2 questions were as good as more questions for detecting depression, 10

Primary prevention

Pravastatin cost was £20 375 (U.S. \$34 360) per life-year gained for primary prevention among men with increased risk for cardiovascular disease, 80

Review: Patient education and counseling increase preventive behaviors, 68

Q

Questionnaires

2 questions were as good as more questions for detecting depression, 10

R

Radioiodine therapy

Radioiodine plus prednisone prevented development or progression of ophthalmopathy in patients with Graves hyperthyroidism, 71

Randomised Intervention Treatment of Angina (RITA-2)

Coronary angioplasty led to more nonfatal MIs than medical therapy in patients with coronary artery disease, 36

Recurrence

Eradication of *Helicobacter pylori* before NSAID treatment prevented ulcers, 44

Review: Azathioprine maintains remission in Crohn disease, but withdrawal rates are higher, 33

Rehabilitation

Early discharge to community care was as effective as conventional care for stroke, 58

Reminder systems

Review: Prompted follow-up by specialists and GPs produces similar outcomes for diabetes mellitus, 57

Review: Telephone and computer communication improves clinical health care process and patient outcome, 22

Respiratory insufficiency

Review: Noninvasive positive-pressure ventilation reduces mortality and need for intubation in the intensive care unit, 39

Rifampin

A catheter coated with minocycline and rifampin reduced bloodstream infections, 41

Roxithromycin

In a pilot study, roxithromycin reduced 1 of 10 end points of ischemic cardiac disease, 35

S

Seizures

Review: Benzodiazepines are more effective than neuroleptics in reducing delirium and seizures in alcohol withdrawal, 8

Septicemia

Antiseptic catheters reduced catheter-related infections and were well tolerated, 40

A catheter coated with minocycline and rifampin reduced bloodstream infections, 4

Serotonin uptake inhibitors

Review: Selective serotonin reuptake inhibitors are effective for mixed chronic pain, 3

Sick-sinus syndrome

Atrial pacing was better than ventricular pacing for the sick-sinus syndrome, 62

Silver sulfadiazine

Antiseptic catheters reduced catheter-related infections and were well tolerated, 40

Smoking

Review: Smoking is associated with increased hip fracture in postmenopausal women, 49

Switching from cigarettes to pipes or cigars reduced smoking-related mortality, 18

Somatostatin

Review: Somatostatin or octreotide reduces transfusion needs but not other outcomes in acute bleeding esophageal varices, 9

Streptokinase

Older age, lighter weight, female sex, African ancestry, and invasive procedures were associated with bleeding after thrombolytic therapy, 17

Surgery

Review: Low-molecular-weight heparin is at least as effective as unfractionated heparin for preventing thrombosis and embolism after surgery, 5

T

Tacrine

Review: Tacrine has limited efficacy for Alzheimer disease, 46

Telephone

Review: Telephone and computer communication improves clinical health care process and patient outcome, 22

Thromboembolism

Atrial pacing was better than ventricular pacing for the sick-sinus syndrome, 62

Review: Factor V Leiden mutation increases risk for thromboembolism, 52

Review: Low-molecular-weight heparin is at least as effective as unfractionated heparin for preventing thrombosis and embolism after surgery, 5

Thrombolytic therapy

Benefit from thrombolysis in acute MI was predicted in the emergency department, 25

Invasive treatment reduced reinfarctions and admissions for unstable angina in patients with ischemia after MI, 4

Older age, lighter weight, female sex, African ancestry, and invasive procedures were associated with bleeding after thrombolytic therapy, 17

Review: Early angioplasty improves outcomes for patients with myocardial infarction, 59

Review: Effectiveness of local opinion leaders varies, 23

Thrombophlebitis

Review: Low-molecular-weight heparin is at least as effective as unfractionated heparin for preventing thrombosis and embolism after surgery, 5

U

Ultrasonography

Serial ultrasonography did not predict stroke in asymptomatic carotid artery disease, 12

V

Varices, esophageal and gastric

Review: Somatostatin or octreotide reduces transfusion needs but not other outcomes in acute bleeding esophageal varices, 9

Vascular disease

Elevated total homocysteine levels increased the risk for vascular disease, 21

Ventilators, mechanical

Review: Noninvasive positive-pressure ventilation reduces mortality and need for intubation in the intensive care unit, 39

Ventricular dysfunction

Review: Subtle clinical findings can detect left-sided heart failure in adults, 11

Ventricular fibrillation

Implantable cardioverter-defibrillators reduced mortality in patients resuscitated from ventricular arrhythmias, 60

Implantable cardioverter-defibrillators used at the time of bypass surgery did not improve survival among high-risk patients, 61

Viral load

HIV-1 RNA levels and CD4+ cell counts predicted disease progression in HIV infection, 13

Vitamin D

Calcium and vitamin D reduced nonvertebral fractures and bone loss, 47

Vitamin E

Vitamin E and β -carotene were not effective in men at high risk for coronary events, 7

W

Women

Dietary intake of monounsaturated and polyunsaturated fats reduced risk for coronary disease in women, 75

Extending screening mammography to include younger women minimally increased life expectancy with large increases in cost, 79

HRT was not associated with increased breast cancer in women with a positive family history of breast cancer, 74

Postmenopausal hormone use was associated with decreased total mortality, 14

Review: HRT is not associated with cardiovascular events or cancer in postmenopausal women in recent clinical trials, 48

Review: Recent HRT increases the risk for breast cancer, 73

Review: Smoking is associated with increased hip fracture in postmenopausal women, 49

Z

Zidovudine

Indinavir plus zidovudine and lamivudine reduced HIV RNA levels in most HIV-infected persons with previous antiretroviral treatment, 42

Indinavir plus zidovudine and lamivudine reduced the progression to AIDS in patients with previous zidovudine therapy, 43

